

ORTHOGONAL POLYNOMIALS IN SEVERAL VARIABLES. I

T. CONSTANTINESCU

ABSTRACT. In this paper we introduce and discuss some classes of orthogonal polynomials in several non-commuting variables. The emphasis is on a non-commutative version of the orthogonal polynomials on the real line. We introduce recurrence equations for these polynomials, Christoffel-Darboux formulas, and Jacobi type matrices.

1. INTRODUCTION

Orthogonal polynomials in several variables are known for long time, see for instance [8], but their theory is less developed than in the one variable case. The commutative case is also studied more intensively (see, for instance, [2], [18]), while the studies for the non-commutative case appear to be quite sparse (see [11]).

Our goal is to introduce and study some classes of orthogonal polynomials in several non-commuting variables. In this paper we focus on polynomials that are viewed as analogues of the orthogonal polynomials on the real line. The main topics are: recurrence equations, Szegő kernels, and Jacobi matrices.

The paper is organized as follows. In Section 2 we introduce the main definitions and several examples. Especially, we briefly review a non-commutative version of the Szegő theory of orthogonal polynomials on the unit circle that was sketched in [7]. Section 3 deals with recurrence equations. In Section 4 we introduce a non-commutative Szegő type kernel which is viewed as a sort of reproducing kernel for the Siegel upper half-space. Section 5 deals with Jacobi type matrices associated to the recurrence equations introduced in Section 3.

2. ORTHOGONAL POLYNOMIALS

In this section we introduce the main definitions and briefly discuss several examples. Let $\mathbb{F}_M^+$ be the unital free semigroup on M generators $g_1, \dots, g_M$ with lexicographic order $\prec$. The empty word is the identity element and the length of the word σ is denoted by $|\sigma|$. The length of the empty word is 0.

Let $\mathcal{O}_N^0$ be the algebra of polynomials in $2N$ non-commuting indeterminates $Y_1, \dots, Y_N, Y_{N+1}, \dots, Y_{2N}$ with complex coefficients. Each element $Q \in \mathcal{O}_N^0$ can be uniquely written in the form $Q = \sum_{\sigma \in \mathbb{F}_{2N}^+} c_\sigma Y_\sigma$, with only finitely many $c_\sigma \neq 0$ and $Y_\sigma = Y_{i_1} \dots Y_{i_k}$ for $\sigma = i_1 \dots i_k \in \mathbb{F}_{2N}^+$. An involution $\mathcal{I}$ can be introduced on $\mathcal{O}_N^0$ as follows:

$$\mathcal{I}(Y_k) = Y_{N+k}, \quad k = 1, \dots, N,$$

$$\mathcal{I}(Y_l) = Y_{l-N}, \quad l = N+1, \dots, 2N;$$

on monomials,

$$\mathcal{I}(Y_{i_1} \dots Y_{i_k}) = \mathcal{I}(Y_{i_k}) \dots \mathcal{I}(Y_{i_1}),$$

and finally, if $Q = \sum_{\sigma \in \mathbb{F}_{2N}^+} c_\sigma Y_\sigma$, then $\mathcal{I}(Q) = \sum_{\sigma \in \mathbb{F}_{2N}^+} \bar{c}_\sigma \mathcal{I}(Y_\sigma)$. Thus, $\mathcal{O}_N^0$ is a unital, associative, $*$ -algebra over $\mathbb{C}$.

Let $\mathcal{P}_N^0$ denote the algebra of polynomials in N non-commuting indeterminates $Y_1, \dots, Y_N$ with complex coefficients. Then $\mathcal{P}_N^0$ is a subalgebra of $\mathcal{O}_N^0$. We say that $\mathcal{A} \subset \mathcal{O}_N^0$ is $\mathcal{I}$ -symmetric if $P \in \mathcal{A}$ implies $c\mathcal{I}(P) \in \mathcal{A}$ for some $c \in \mathbb{C} - \{0\}$. We construct an associative algebra $\mathcal{O}_N^0(\mathcal{A})$ as the quotient of $\mathcal{O}_N^0$ by the two-sided ideal $\mathcal{E}(\mathcal{A})$ generated by $\mathcal{A}$. We notice that $\mathcal{O}_N^0(\emptyset) = \mathcal{O}_N^0$. We let $\pi = \pi_{\mathcal{A}} : \mathcal{O}_N^0 \rightarrow \mathcal{O}_N^0(\mathcal{A})$ be the quotient map and since $\mathcal{A}$ is $\mathcal{I}$ -symmetric,

$$(2.1) \quad \mathcal{I}_{\mathcal{A}}(\pi(P)) = \pi(\mathcal{I}(P))$$

gives an involution on $\mathcal{O}_N^0(\mathcal{A})$. We will be interested in linear functionals ϕ on $\mathcal{O}_N^0(\mathcal{A})$ with the property that $\phi(\mathcal{I}_{\mathcal{A}}(\pi(P))\pi(P)) \geq 0$ for all $P \in \mathcal{P}_N^0$. Without loss of generality we will assume that ϕ is unital, $\phi(\pi(1)) = 1$. Such a functional will be called a *positive functional* on $\mathcal{O}_N^0(\mathcal{A})$. The proof of the following result is straightforward and can be omitted.

Lemma 1. *Let ϕ be a positive functional on $\mathcal{O}_N^0(\mathcal{A})$. Then*

- 1) $\phi(\mathcal{I}_{\mathcal{A}}(\pi(P))) = \overline{\phi(\pi(P))}$ for $P \in \mathcal{P}_N^0$.
- 2) $|\phi(\mathcal{I}_{\mathcal{A}}(\pi(P_1))\pi(P_2))|^2 \leq \phi(\mathcal{I}_{\mathcal{A}}(\pi(P_1))\pi(P_1))\phi(\mathcal{I}_{\mathcal{A}}(\pi(P_2))\pi(P_2))$ for $P_1, P_2 \in \mathcal{P}_N^0$.

We now consider the GNS construction associated to ϕ . Thus, we define on $\pi(\mathcal{P}_N^0)$,

$$(2.2) \quad \langle \pi(P_1), \pi(P_2) \rangle_\phi = \phi(\mathcal{I}_{\mathcal{A}}(\pi(P_2))\pi(P_1)),$$

and factor out the subspace $\mathcal{N}_\phi = \{\pi(P) \mid P \in \mathcal{P}_N^0, \langle \pi(P), \pi(P) \rangle_\phi = 0\}$. Completing this quotient with respect to the norm induced by (2.2) we obtain a Hilbert space $\mathcal{H}_\phi$. From now on we will assume that ϕ is strictly positive, that is, $\phi(\mathcal{I}_{\mathcal{A}}(\pi(P))\pi(P)) > 0$ for all $P \in \mathcal{P}_N^0 - \mathcal{E}(\mathcal{A})$, so that $\mathcal{N}_\phi = \{0\}$ and $\pi(\mathcal{P}_N^0)$ can be viewed as a subspace of $\mathcal{H}_\phi$. Let $\mathcal{F} = \{F_\alpha\}_{\alpha \in G}$ be the set of the distinct elements $\pi(Y_\sigma)$, $\sigma \in \mathbb{F}_N^+$. The *index set* G is chosen as follows: the equality on $\{\pi(Y_\sigma) \mid \sigma \in \mathbb{F}_N^+\}$ gives an equivalence relation and choose from each equivalence class the element $\pi(Y_\sigma)$ with the least σ with respect to the lexicographic order. Then $G \subset \mathbb{F}_N^+$ and $\emptyset \in G$. Let $G_n = \{\alpha \in G \mid |\alpha| = n\}$, then $G_0 = \{\emptyset\}$ and $\{G_n\}_{n \geq 0}$ is a partition of G .

Since ϕ is strictly positive it follows that $\mathcal{F}$ is a linearly independent family in $\mathcal{H}_\phi$ and the Gram-Schmidt procedure gives a family $\{\varphi_\alpha\}_{\alpha \in G}$ of elements in $\pi(\mathcal{P}_N^0) \subset \mathcal{O}_N^0(\mathcal{A})$ such that

$$(2.3) \quad \varphi_\alpha = \sum_{\beta \preceq \alpha} a_{\alpha, \beta} F_\beta, \quad a_{\alpha, \alpha} > 0;$$

$$(2.4) \quad \langle \varphi_\alpha, \varphi_\beta \rangle_\phi = \delta_{\alpha,\beta}, \quad \alpha, \beta \in G.$$

The elements φ_α , $\alpha \in G$, will be called the *orthogonal polynomials* associated to ϕ . Typically, the theory of orthogonal polynomials deals with the study of algebraic and asymptotic properties of the orthogonal polynomials associated to strictly positive functionals on $\mathcal{O}_N^0(\mathcal{A})$. An explicit formula for the orthogonal polynomials can be obtained in the same manner as in the classical (one variable) case. Define

$$(2.5) \quad s_{\alpha,\beta} = \phi(\mathcal{I}_{\mathcal{A}}(F_\alpha)F_\beta) = \langle F_\beta, F_\alpha \rangle_\phi, \quad \alpha, \beta \in G,$$

and

$$(2.6) \quad D_\alpha = \det [s_{\alpha',\beta'}]_{\alpha',\beta' \preceq \alpha} > 0, \quad \alpha \in G.$$

We notice that ϕ is a positive functional on $\mathcal{O}_N^0(\mathcal{A})$ if and only if $K_\phi(\alpha, \beta) = s_{\alpha,\beta}$, $\alpha, \beta \in G$, is a positive definite kernel on G . From now on $\tau - 1$ denotes the predecessor of τ with respect to the lexicographic order $\prec$ on $\mathbb{F}_N^+$, while $\sigma + 1$ denotes the successor of σ . The determinant involved in the next result is defined by the same formula as in the scalar case, even though its entries are elements of $\pi(\mathcal{P}_N^0)$.

Theorem 2. *Let $\{\varphi_\alpha\}_{\alpha \in G}$ be the orthogonal polynomials associated to the strictly positive, unital functional ϕ on $\mathcal{O}_N^0(\mathcal{A})$. Then $\varphi_\emptyset = 1$ and for $\emptyset \prec \alpha$,*

$$(2.7) \quad \varphi_\alpha = \frac{1}{\sqrt{D_{\alpha-1}D_\alpha}} \det \begin{bmatrix} [s_{\alpha',\beta'}]_{\alpha' \prec \alpha; \beta' \preceq \alpha} \\ F_\emptyset \quad \dots \quad F_\alpha \end{bmatrix}.$$

Proof. The proof is similar to the classical one. Thus, we deduce from the orthogonality condition (2.4) that $\langle \varphi_\alpha, F_{\beta'} \rangle_\phi = 0$ for $\emptyset \preceq \beta' \prec \alpha$, which implies that $\sum_{\beta \preceq \alpha} a_{\alpha,\beta} s_{\beta',\beta} = 0$ for $\emptyset \preceq \beta' \prec \alpha$. Since the coefficients of the linear system

$$\begin{cases} \sum_{\beta \preceq \alpha} a_{\alpha,\beta} s_{\beta',\beta} = 0, & \emptyset \preceq \beta' \prec \alpha, \\ \sum_{\beta \preceq \alpha} a_{\alpha,\beta} F_\beta = \varphi_\alpha, \end{cases}$$

with unknowns $a_{\alpha,\beta}$ are complex numbers except for those of the last equation which are in $\pi(\mathcal{P}_N^0)$, Cramer's rule still holds in a form that gives

$$\varphi_\alpha = \frac{a_{\alpha,\alpha}}{D_{\alpha-1}} \det \begin{bmatrix} [s_{\alpha',\beta'}]_{\alpha' \prec \alpha; \beta' \preceq \alpha} \\ F_\emptyset \quad \dots \quad F_\alpha \end{bmatrix}.$$

Next we notice that

$$\langle \det \begin{bmatrix} [s_{\alpha',\beta'}]_{\alpha' \prec \alpha; \beta' \preceq \alpha} \\ F_\emptyset \quad \dots \quad F_\alpha \end{bmatrix}, F_\alpha \rangle_\phi = D_\alpha,$$

and since $F_\alpha = \frac{1}{a_{\alpha,\alpha}}\varphi_\alpha + \sum_{\beta \prec \alpha} c_\beta F_\beta$ for some complex coefficients c_β , $\beta \prec \alpha$, we deduce

$$D_\alpha = \langle \frac{D_{\alpha-1}}{a_{\alpha,\alpha}}\varphi_\alpha, \frac{1}{a_{\alpha,\alpha}}\varphi_\alpha + \sum_{\beta \prec \alpha} c_\beta F_\beta \rangle_\phi = \frac{D_{\alpha-1}}{a_{\alpha,\alpha}^2},$$

so that

$$\frac{1}{a_{\alpha,\alpha}^2} = \frac{D_\alpha}{D_{\alpha-1}},$$

which gives (2.7). $\square$

Usually, the representation (2.7) is not very useful for the actual computation of the orthogonal polynomials. Instead, recurrence equations are obtained for each particular case of interest. We consider several such examples.

2.1. Orthogonal polynomials in one variable. Assume $N = 1$. Then a linear functional ϕ on $\mathcal{O}_1^0$ is positive if and only if $K_\phi(n, m) = \phi(\mathcal{I}(Y_1^n)Y_1^m)$, $n, m \in \mathbb{N}$, is a positive definite kernel on $\mathbb{N}$. One simple example can be obtained by taking $\mathcal{A}_{\mathbb{T}} = \{1 - \mathcal{I}(Y_1)Y_1\}$. In this case, ϕ is a positive functional on $\mathcal{O}_1^0(\mathcal{A}_{\mathbb{T}})$ if and only if the kernel K_ϕ is positive definite and satisfy the Toeplitz condition, that is

$$K_\phi(n + k, m + k) = K_\phi(n, m), \quad m, n, k \in \mathbb{N}.$$

This shows that the orthogonal polynomials on $\mathcal{O}_1^0(\mathcal{A}_{\mathbb{T}})$ are the orthogonal polynomials on the unit circle ([17]). The index set is $\mathbb{N}$ in this case, and define $\gamma_n = -a_{n,n}\varphi_n(0)$ for $n \geq 1$. Then $|\gamma_n| < 1$ and set $d_n = (1 - |\gamma_n|^2)^{1/2}$. The orthogonal polynomials on the unit circle satisfy the following basic recurrence equations (see [17]):

$$(2.8) \quad \varphi_{n+1}(z) = \frac{1}{d_{n+1}}(z\varphi_n(z) - \gamma_{n+1}\varphi_n^\sharp(z)),$$

where $\varphi_0^\sharp(z) = 1$ and for $n > 0$,

$$(2.9) \quad \varphi_{n+1}^\sharp(z) = \frac{1}{d_{n+1}}(-\overline{\gamma}_{n+1}z\varphi_n(z) + \varphi_n^\sharp(z)).$$

Another example is given by $\mathcal{A}_{\mathbb{R}} = \{Y_1 - \mathcal{I}(Y_1)\}$. The index set is still $\mathbb{N}$, but this time, ϕ is a positive functional on $\mathcal{O}_1^0(\mathcal{A}_{\mathbb{R}})$ if and only if the kernel K_ϕ is positive definite and has the Hankel property, that is

$$K_\phi(n, m + k) = K_\phi(n + k, m), \quad m, n, k \in \mathbb{N}.$$

This shows that the orthogonal polynomials on $\mathcal{O}_1^0(\mathcal{A}_{\mathbb{R}})$ are the orthogonal polynomials on the real line. In this case one obtains a basic three-terms recurrence equation (see [17]):

$$(2.10) \quad x\varphi_n(x) = b_{n+1}\varphi_{n+1}(x) + a_n\varphi_n(x) + b_n\varphi_{n-1}(x),$$

with initial conditions $\varphi_0(x) = 1$ and $\varphi_{-1}(x) = 0$.

2.2. *Examples in several variables.* In this case, there is a large number of interesting examples. Here we mention just one. More examples will be considered in the next sections. Let

$$\mathcal{A}_1 = \{1 - \mathcal{I}(Y_k)Y_k \mid k = 1, \dots, N\} \cup \{\mathcal{I}(Y_k)Y_l \mid k, l = 1, \dots, N, k \neq l\}.$$

Then ϕ is a positive functional on $\mathcal{O}_N^0(\mathcal{A}_1)$ if and only if K_ϕ is a positive definite kernel that obeys the rules:

$$(2.11) \quad K_{\tilde{\phi}}(\tau\sigma, \tau\sigma') = K_{\tilde{\phi}}(\sigma, \sigma'), \quad \tau, \sigma, \sigma' \in \mathbb{F}_N^+,$$

$$(2.12) \quad K_{\tilde{\phi}}(\sigma, \tau) = 0 \quad \text{if there is no } \alpha \in \mathbb{F}_N^+ \text{ such that } \sigma = \alpha\tau \text{ or } \tau = \alpha\sigma.$$

Such type of kernels appeared in the study of some classes of stochastic processes indexed by nodes on a tree (see, for instance, [3]). In this case the index set is $\mathbb{F}_N^+$. We also need to introduce

$$(2.13) \quad D_{1,\sigma} = \det [K_{\sigma',\tau'}]_{\emptyset \prec \sigma', \tau' \preceq \sigma} > 0.$$

Define $\gamma_\sigma = -\sqrt{\frac{D_\sigma}{D_{1,\sigma}}}a_{\sigma,\emptyset}$ and notice that $|\gamma_\sigma| < 1$. Then we can define $d_\sigma = (1 - |\gamma_\sigma|^2)^{1/2}$ and the orthogonal polynomials associated to $\mathcal{O}_N^0(\mathcal{A}_1)$ obey the recurrence equations (see [7]):

$$(2.14) \quad \varphi_{k\sigma} = \frac{1}{d_{k\sigma}}(Y_k\varphi_\sigma - \gamma_{k\sigma}\varphi_{k\sigma-1}^\sharp), \quad k = 1, \dots, N, \quad \sigma \in \mathbb{F}_N^+,$$

where $\varphi_\emptyset^\sharp = 1$ and for $k \in \{1, \dots, N\}$, $\sigma \in \mathbb{F}_N^+$,

$$(2.15) \quad \varphi_{k\sigma}^\sharp = \frac{1}{d_{k\sigma}}(-\overline{\gamma}_{k\sigma}Y_k\varphi_\sigma + \varphi_{k\sigma-1}^\sharp).$$

3. RECURRENCE EQUATIONS

In this section we consider some algebraic properties of the orthogonal polynomials of $\mathcal{O}_N^0(\mathcal{A}_2)$, where $\mathcal{A}_2 = \{Y_k - \mathcal{I}(Y_k) \mid k = 1, \dots, N\}$. If we take $\mathcal{A}_2' = \mathcal{A}_2 \cup \{Y_kY_l - Y_lY_k \mid k, l = 1, \dots, N\}$, then $\pi_{\mathcal{A}_2'}(\mathcal{P}_N^0)$ is isomorphic to the symmetric algebra of $\mathbb{C}^N$ and the orthogonal polynomials correspond to the orthogonal polynomials of several commuting variables, see [18]. Since $\pi_{\mathcal{A}_2'}(\mathcal{P}_N^0)$ is a quotient of $\pi_{\mathcal{A}_2}(\mathcal{P}_N^0) = \mathcal{P}_N^0$, we expect that results for $\mathcal{P}_N^0$ would give corresponding results for $\pi_{\mathcal{A}_2'}(\mathcal{P}_N^0)$ by standard (symmetrization) techniques, see [14]. Going in the opposite direction, we expect to deduce generalizations of results on $\pi_{\mathcal{A}_2'}(\mathcal{P}_N^0)$ to similar results for $\mathcal{P}_N^0$. We illustrate this remark by obtaining a three-term recurrence relation for the orthogonal polynomials in $\mathcal{O}_N^0(\mathcal{A}_2)$.

Let ϕ be a strictly positive functional on $\mathcal{O}_N^0(\mathcal{A}_2)$. Since $\pi_{\mathcal{A}_2}(\mathcal{P}_N^0) = \mathcal{P}_N^0$, the index set G is $\mathbb{F}_N^+$ and G_n is the set of words in $\mathbb{F}_N^+$ of length n . Let $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ be the orthogonal

polynomials associated to ϕ . The matrix-vector notation in [12] is easily adapted to $\mathcal{P}_N^0$, by setting $\Phi_n = [\varphi_\sigma]_{|\sigma|=n}$, $n \geq 0$. We can show that (2.10) extends to $\mathcal{O}_N^0(\mathcal{A}_2)$.

Theorem 3. *The orthogonal polynomials on $\mathcal{O}_N^0(\mathcal{A}_2)$ obey the following recurrence relations: for $k = 1, \dots, N$,*

$$(3.1) \quad Y_k \Phi_0 = \Phi_1 B_{0,k} + \Phi_0 A_{0,k},$$

and for $k = 1, \dots, N$ and $n \geq 1$,

$$(3.2) \quad Y_k \Phi_n = \Phi_{n+1} B_{n,k} + \Phi_n A_{n,k} + \Phi_{n-1} B_{n-1,k}^*.$$

Proof. The proof is most like in the classical, one-dimensional case. Thus, we can write

$$Y_k \varphi_\sigma = \sum_{\tau \preceq k\sigma} c_\tau^{k\sigma} \varphi_\tau, \quad k \in \{1, \dots, N\}, \quad \sigma \in \mathbb{F}_N^+,$$

where the coefficients $c_\tau^{k\sigma}$ are calculated by the formula $c_\tau^{k\sigma} = \langle Y_k \varphi_\sigma, \varphi_\tau \rangle_\phi$. We notice that for any $P, Q \in \mathcal{P}_N^0$,

$$\begin{aligned} \langle Y_k P, Q \rangle_\phi &= \phi(\mathcal{I}(Q) Y_k P) \\ (3.3) \quad &= \phi(\mathcal{I}(Q) \mathcal{I}(Y_k) P) \\ &= \langle P, Y_k Q \rangle_\phi. \end{aligned}$$

[More generally, we have $s_{\alpha\sigma,\tau} = s_{\sigma, I(\alpha)\tau}$ for $\alpha, \sigma, \tau \in \mathbb{F}_N^+$, where I denotes the involution on $\mathbb{F}_N^+$ given by $I(i_1 \dots i_k) = i_k \dots i_1$]. In particular, for $|\tau| \leq |\sigma| - 2$,

$$c_\tau^{k\sigma} = \langle \varphi_\sigma, Y_k \varphi_\tau \rangle_\phi = 0,$$

while for the remaining values of $\tau \preceq k\sigma$,

$$(3.4) \quad c_\tau^{k\sigma} = \langle \varphi_\sigma, Y_k \varphi_\tau \rangle_\phi = \overline{\langle Y_k \varphi_\tau, \varphi_\sigma \rangle_\phi} = \bar{c}_\sigma^{k\tau}.$$

We deduce that for $k = 1, \dots, N$,

$$(3.5) \quad Y_k \Phi_0 = \Phi_1 B_{0,k} + \Phi_0 A_{0,k},$$

while for $k = 1, \dots, N$ and $n \geq 1$,

$$(3.6) \quad Y_k \Phi_n = \Phi_{n+1} B_{n,k} + \Phi_n A_{n,k} + \Phi_{n-1} C_{n,k}.$$

Let $X = [x_{ij}]$ be a given matrix. We will use the following notation: first, $\mathcal{I}(X) = [\mathcal{I}(x_{ji})]$ and then $\phi(X) = [\phi(x_{ij})]$. We deduce from (3.5), (3.6), and (2.4) that

$$(3.7) \quad B_{n,k}^* = \phi(\mathcal{I}(\Phi_n) Y_k \Phi_{n+1}),$$

$$(3.8) \quad A_{n,k}^* = \phi(\mathcal{I}(\Phi_n) Y_k \Phi_n),$$

and

$$(3.9) \quad C_{n,k}^* = \phi(\mathcal{I}(\Phi_n) Y_k \Phi_{n-1}) = \phi(\mathcal{I}(\Phi_{n-1}) Y_k \Phi_n)^* = B_{n-1,k}.$$

□

We notice that (3.3) implies that $A_{n,k}^* = A_{n,k}$ and if we define $B_n = \begin{bmatrix} B_{n,1} & \dots & B_{n,N} \end{bmatrix}$, $n \geq 0$, then B_n is an $N^{n+1} \times N^{n+1}$ upper triangular invertible matrix. Under these conditions we can prove a converse of Theorem 3. This appears as a Favard type result and gives a construction of strictly positive functionals on $\mathcal{O}_N^0(\mathcal{A}_2)$.

Theorem 4. *Let $\varphi_\sigma = \sum_{\tau \preceq \sigma} a_{\sigma,\tau} Y_\tau$, $\sigma \in \mathbb{F}_N^+$, be elements in $\mathcal{P}_N^0$ such that $\varphi_\emptyset = 1$ and $a_{\sigma,\sigma} > 0$. Assume that there exist families $\{A_{n,k} \mid n \geq 0, k = 1, \dots, N\}$, $\{B_{n,k} \mid n \geq 0, k = 1, \dots, N\}$ of matrices such that $A_{n,k}^* = A_{n,k}$ for all $n \geq 0$ and $k = 1, \dots, N$, $B_n = \begin{bmatrix} B_{n,1} & \dots & B_{n,N} \end{bmatrix}$ is an upper triangular invertible matrix for each $n \geq 0$, for $k = 1, \dots, N$,*

$$(3.10) \quad Y_k \varphi_\emptyset = [\varphi_\sigma]_{|\sigma|=1} B_{0,k} + \varphi_\emptyset A_{0,k},$$

and for $k = 1, \dots, N$, $n \geq 1$,

$$(3.11) \quad Y_k [\varphi_\sigma]_{|\sigma|=n} = [\varphi_\sigma]_{|\sigma|=n+1}^T B_{n,k} + [\varphi_\sigma]_{|\sigma|=n}^T A_{n,k} + [\varphi_\sigma]_{|\sigma|=n-1}^T B_{n-1,k}^*.$$

Then there exists a strictly positive functional ϕ on $\mathcal{O}_N^0(\mathcal{A}_2)$ such that $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ is the family of orthogonal polynomials associated to ϕ .

Proof. Since B_n , $n \geq 0$, are invertible matrices, it follows that $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ is a linearly independent family in $\mathcal{P}_N^0$ and formula (2.4) suggests to define:

$$(3.12) \quad \phi(1) = 1 \quad \text{and} \quad \phi(\varphi_\sigma) = 0 \quad \text{for} \quad \sigma \in \mathbb{F}_N^+ - \{\emptyset\}.$$

These relations uniquely determine a linear functional on $\mathcal{O}_N^0(\mathcal{A}_2) = \mathcal{P}_N^0$. We will use again the matrix-vector notation, $\Phi_n = [\varphi_\sigma]_{|\sigma|=n}$, $n \geq 0$. Also, set $\Phi_{-1} = 0$ and $B_{-1,k} = 0$. Then (3.10) can be included in (3.11) for $n = 0$.

We prove that

$$(3.13) \quad \phi(\mathcal{I}(\Phi_n)\Phi_m) = \begin{cases} 0, & n \neq m; \\ I, & n = m, \end{cases}$$

where 0 and I denote the zero, respectively the identity matrix of a suitable dimension. This will imply that ϕ is a strictly positive functional on $\mathcal{P}_N^0$ and that the orthogonal polynomials associated to ϕ are precisely φ_σ , $\sigma \in \mathbb{F}_N^+$.

We prove (3.13) by induction on m . For $m = 0$,

$$\begin{aligned} \phi(\mathcal{I}(\Phi_n)\Phi_0) &= \Phi(\mathcal{I}(\Phi_0)\Phi_n)^* \\ &= [\phi(\varphi_\sigma)]_{|\sigma|=n}^*, \end{aligned}$$

so that, by (3.12), $\phi(\mathcal{I}(\Phi_0)\Phi_0) = 1$ and $\phi(\mathcal{I}(\Phi_n)\Phi_0) = 0$ for $n > 0$. Assume (3.13) holds for $n \geq 0$ and $k \leq m$. Then $\phi(\mathcal{I}(\Phi_n)\Phi_{m+1}) = 0$ for $n \leq m$ by the induction hypothesis. We deduce from (3.11) that

$$[Y_1\Phi_l \dots Y_N\Phi_l] = \Phi_{l+1}B_l + \Phi_l A_l + \Phi_{l-1}C_l,$$

where $A_l = [A_{l,1} \dots A_{l,N}]$ and $C_l = [B_{l-1,1}^* \dots B_{l-1,N}^*]$, hence

$$\begin{aligned} \Phi_{l+1} &= [Y_1\Phi_l \dots Y_N\Phi_l] B_l^{-1} - \Phi_l A_l B_l^{-1} - \Phi_{l-1} C_l B_l^{-1} \\ &= \sum_{j=1}^N Y_j \Phi_l D_{l,j} - \Phi_l E_l - \Phi_{l-1} F_l, \end{aligned}$$

where $B_l^{-1} = [D_{l,1} \dots D_{l,N}]^T$, T denoting the matrix transpose, $E_l = A_l B_l^{-1}$, and $F_l = C_l B_l^{-1}$. Using the induction hypothesis, the previous formula for $l = m$, and (3.11), we deduce

$$\begin{aligned} \phi(\mathcal{I}(\Phi_{m+1})\Phi_{m+1}) &= \sum_{j=1}^N D_{m,j}^* \phi(\mathcal{I}(\Phi_m)Y_j\Phi_{m+1}) \\ &\quad - E_m^* \phi(\mathcal{I}(\Phi_m)\Phi_{m+1}) - F_m^* \phi(\mathcal{I}(\Phi_{m-1})\Phi_{m+1}) \\ &= \sum_{j=1}^N D_{m,j}^* \phi(\mathcal{I}(Y_j\Phi_m)\Phi_{m+1}) \\ &= \sum_{j=1}^N D_{m,j}^* B_{m,j}^* \phi(\mathcal{I}(\Phi_{m+1})\Phi_{m+1}) \\ &\quad + \sum_{j=1}^N D_{m,j}^* B_{m,j}^* \phi(\mathcal{I}(\Phi_m)\Phi_{m+1}) \\ &\quad + \sum_{j=1}^N D_{m,j}^* A_{m-1,j} \phi(\mathcal{I}(\Phi_{m-1})\Phi_{m+1}) \\ &= \sum_{j=1}^N D_{m,j}^* B_{m,j}^* = I. \end{aligned}$$

Similar calculations show that $\phi(\mathcal{I}(\Phi_l)\Phi_{m+1}) = 0$ for $l > m + 1$ and the proof is concluded. $\square$

4. CHRISTOFFEL-DARBOUX FORMULA

In the classical theory orthogonal polynomials are evaluated at points in some suitable domains. For instance, for orthogonal polynomials on the unit circle the domain is the open unit disk and the polynomials are closely related to the theory of analytic functions on that domain. A key to this connection can be considered to be the Szegő kernel $K_S(z, w) = \frac{1}{1-z\bar{w}}$. A similar relation is established between orthogonal polynomials on the real line and function theory on the upper half plane. In this section we suggest an extension of these connections to several non-commuting variables. The analogue of the unit disk was already dealt with in [6], [7], and we begin by briefly reviewing that construction.

Let $\mathcal{E}$ be an infinite-dimensional Hilbert space and by $\mathcal{L}(\mathcal{E})$ we denote the set of bounded linear operators on $\mathcal{E}$. The N -dimensional unit ball of $\mathcal{E}$ is defined by

$$\mathcal{B}_N(\mathcal{E}) = \{Z = \begin{pmatrix} Z_1 & \dots & Z_N \end{pmatrix} \mid (Z \mid Z) < I_{\mathcal{E}}\},$$

where $I_{\mathcal{E}}$ is the identity on $\mathcal{E}$ and for $Z = \begin{pmatrix} Z_1 & \dots & Z_N \end{pmatrix}$ and $Z' = \begin{pmatrix} Z'_1 & \dots & Z'_N \end{pmatrix}$ in $\mathcal{L}(\mathcal{E})^N$,

$$(4.1) \quad (Z \mid Z') = \sum_{k=1}^N Z_k(Z'_k)^*.$$

A family of Hilbert spaces is associated to the Hilbert space $\mathcal{E}$ as follows: $\mathcal{E}_0 = \mathcal{E}$ and for $k \geq 1$,

$$\mathcal{E}_k = \underbrace{\mathcal{E}_{k-1} \oplus \dots \oplus \mathcal{E}_{k-1}}_{N \text{ terms}} = \mathcal{E}_{k-1}^{\oplus N}.$$

For $\mathcal{E} = \mathbb{C}$ we deduce $\mathbb{C}_k = (\mathbb{C}^N)^{\otimes k}$, the k -fold tensor product of $\mathbb{C}^N$ with itself, therefore $\oplus_{k \geq 0} \mathbb{C}_k$ is the Fock space $\mathcal{F}(\mathbb{C}^N)$ associated to $\mathbb{C}^N$. We also deduce that $\oplus_{k \geq 0} \mathcal{E}_k$ is isomorphic to $\mathcal{F}(\mathbb{C}^N) \otimes \mathcal{E}$.

For $Z \in \mathcal{B}_N(\mathcal{E})$ we define $E(Z) = [Z_{\sigma}]_{|\sigma|=0}^{\infty}$ and notice that $E(Z)$ is a bounded linear operator from $\oplus_{k \geq 0} \mathcal{E}_k$ into $\mathcal{E}$. The *Szegő kernel* for $\mathcal{B}_N(\mathcal{E})$ is defined by the formula

$$K_{\mathcal{B}}(Z, Z') = E(Z)E(Z')^*, \quad Z, Z' \in \mathcal{B}_N(\mathcal{E}).$$

The following result describes some of the basic properties of $K_{\mathcal{B}}$.

Lemma 5. (a) $K_{\mathcal{B}}$ is a positive definite kernel on $\mathcal{B}_N(\mathcal{E})$.

(b) The set $\{E(Z)^*\mathcal{E} \mid Z \in \mathcal{B}_N(\mathcal{E})\}$ is total in $\oplus_{k \geq 0} \mathcal{E}_k$.

(c) For any $T \in \mathcal{L}(\mathcal{E})$ and $Z, Z' \in \mathcal{B}_N(\mathcal{E})$,

$$E(Z)(T - \sum_{k=1}^N Z_k T(Z'_k)^*)^{\oplus \infty} E(Z')^* = T,$$

where $(T - \sum_{k=1}^N Z_k T(Z'_k)^*)^{\oplus \infty}$ is the diagonal operator in $\mathcal{L}(\oplus_{k \geq 0} \mathcal{E}_k)$ with diagonal entry $T - \sum_{k=1}^N Z_k T(Z'_k)^*$.

Proof. The proof of this result can be found in [7]. Thus, (a) and (c) are quite straightforward. The most interesting is (b). Its proof depends on the assumption the $\mathcal{E}$ is infinite dimensional. In fact, for $\mathcal{E} = \mathbb{C}$, the result is not true, since the set $\{E(Z)^*\mathbb{C} \mid Z \in \mathcal{B}_N(\mathbb{C})\}$ is total in the symmetric Fock space, see [1]. For this reason and sake of completeness, we sketch the proof of (b) here. Let $f = \{f_{\sigma}\}_{\sigma \in \mathbb{F}_N^+}$ be an element of $\oplus_{k \geq 0} \mathcal{E}_k$ orthogonal to the linear span of $\{E(Z)^*\mathcal{E} \mid Z \in \mathcal{B}_N(\mathcal{E})\}$. Taking $Z = 0$, we deduce that $f_{\emptyset} = 0$. Next, we claim that for each $\sigma \in \mathbb{F}_N^+ - \{\emptyset\}$ there exist

$$Z_l = (Z_1^l, \dots, Z_N^l) \in \mathcal{B}_N(\mathcal{E}), \quad l = 1, \dots, 2|\sigma|,$$

such that

$$\text{range} \begin{bmatrix} Z_{\sigma}^{*1} & \dots & Z_{\sigma}^{*2|\sigma|} \end{bmatrix} = \mathcal{E},$$

and

$$Z_\tau^l = 0 \quad \text{for all } \tau \neq \sigma, \quad |\tau| \geq |\sigma|, \quad l = 1, \dots, 2|\sigma|.$$

Once this claim is proved, a simple inductive argument gives $f = 0$, so $\{E(Z)^*\mathcal{E} \mid Z \in \mathcal{B}_N(\mathcal{E})\}$ is total in $\oplus_{k \geq 0} \mathcal{E}_k$. In order to prove the claim we need the following construction.

Let $\{e_{ij}^n\}_{i,j=1}^n$ be the matrix units of the algebra M_n of $n \times n$ matrices. Each e_{ij}^n is an $n \times n$ matrix consisting of 1 in the (i, j) th entry and zeros elsewhere. For a Hilbert space $\mathcal{E}_1$ we define $E_{ij}^n = e_{ij}^n \otimes I_{\mathcal{E}_1}$ and we notice that

$$(4.2) \quad E_{ij}^n E_{kl}^n = \delta_{jk} E_{il}^n, \quad E_{ji}^{*n} = E_{ij}^n.$$

Let $\sigma = i_1 \dots i_k$ so that $\mathcal{E} = \mathcal{E}_1^{\oplus 2|\sigma|}$ for some Hilbert space $\mathcal{E}_1$ (here we essentially use the assumption that $\mathcal{E}$ is of infinite dimension). Also, for $s = 1, \dots, N$, we define

$$J_s = \{l \in \{1, \dots, k\} \mid i_{k+1-l} = s\}$$

and

$$Z_s^{*p} = \frac{1}{\sqrt{2}} \sum_{r \in J_s} E_{r+p-1, r+p}^{2|\sigma|}, \quad s = 1, \dots, N, \quad p = 1, \dots, |\sigma|.$$

We can show that for each $p \in \{1, \dots, |\sigma|\}$,

$$(4.3) \quad Z_\sigma^{*p} = \frac{1}{\sqrt{2^k}} E_{p, k+p}^{2|\sigma|},$$

$$(4.4) \quad Z_\tau^p = 0 \quad \text{for } \tau \neq \sigma, \quad |\tau| \geq |\sigma|.$$

Using (4.2), we deduce

$$\begin{aligned} \sum_{s=1}^N Z_s^p Z_s^{*p} &= \frac{1}{2} \sum_{s=1}^N \sum_{r \in J_s} E_{r+p, r+p-1}^{2|\sigma|} E_{r+p-1, r+p}^{2|\sigma|} \\ &= \frac{1}{2} \sum_{s=1}^N \sum_{r \in J_s} E_{r+p, r+p}^{2|\sigma|} \\ &= \frac{1}{2} \sum_{r=1}^k E_{r+p, r+p}^{2|\sigma|} < I, \end{aligned}$$

hence $Z^p \in \mathcal{B}_N(\mathcal{E})$ for each $p = 1, \dots, |\sigma|$. For each word $\tau = j_1 \dots j_k \in \mathbb{F}_N^+ - \{\emptyset\}$ we deduce by induction that

$$(4.5) \quad Z_{j_k}^{*p} \dots Z_{j_1}^{*p} = \frac{1}{\sqrt{2^k}} \sum_{r \in A_\tau} E_{r+p-1, r+p+k-1}^{2|\sigma|},$$

where $A_\tau = \cap_{p=0}^{k-1} (J_{j_{k-p}} - p) \subset \{1, \dots, N\}$ and $J_{j_{k-p}} - p = \{l - p \mid l \in J_{j_{k-p}}\}$.

We show that $A_\sigma = \{1\}$ and $A_\tau = \emptyset$ for $\tau \neq \sigma$. Let $q \in A_\tau$. Therefore, for any $p \in \{0, \dots, k-1\}$ we must have $q + p \in J_{j_{k-p}}$ or $i_{k+1-q-p} = j_{k-p}$. For $p = k-1$ we deduce $j_1 = i_{2-q}$ and since $2-q \geq 1$, it follows that $q \leq 1$. Also $q \geq 1$, therefore the only element that can be in A_τ is $q = 1$, in which case we must have $\tau = \sigma$. Since $l \in J_{i_{k+1-l}}$ for each $l = 1, \dots, k-1$, hence $A_\sigma = \{1\}$ and $A_\tau = \emptyset$ for $\tau \neq \sigma$. Formula (4.5) implies

(4.3). In a similar manner we can construct a family Z^p , $p = |\sigma| + 1, \dots, 2|\sigma|$, such that

$$Z_\sigma^{*p} = \frac{1}{\sqrt{2^k}} E_{p+k,p}^{2|\sigma|},$$

and

$$Z_\tau^p = 0 \quad \text{for } \tau \neq \sigma, \quad |\tau| \geq |\sigma|.$$

Thus, for $s = 1, \dots, N$, we define

$$K_s = \{l \in \{1, \dots, k\} \mid i_k = s\}$$

and

$$Z_s^{*p} = \frac{1}{\sqrt{2}} \sum_{r \in K_s} E_{r+p-k, r+p-k-1}^{2|\sigma|}, \quad s = 1, \dots, N, \quad p = |\sigma| + 1, \dots, 2|\sigma|.$$

Now,

$$\begin{bmatrix} Z_\sigma^{*1} & \dots & Z_\sigma^{*2|\sigma|} \end{bmatrix} = \frac{1}{\sqrt{2^k}} \begin{bmatrix} E_{1,k+1}^{2|\sigma|} & \dots & E_{k,2k}^{2|\sigma|} & E_{k+1,1}^{2|\sigma|} & \dots & E_{2k,k}^{2|\sigma|} \end{bmatrix},$$

whose range is $\mathcal{E}$. This concludes the proof. $\square$

Define the vector space

$$\mathcal{R} = \{r_f : \mathcal{B}_N(\mathcal{E}) \rightarrow \mathcal{E} \mid r_f(Z) = E(Z)f, \quad f \in \oplus_{k \geq 0} \mathcal{E}_k\},$$

and consider the map $U : \oplus_{k \geq 0} \mathcal{E}_k \rightarrow \mathcal{R}$ defined by $Uf = r_f$. This map is linear and bijective (by Lemma 5 (b)), so we can define on $\mathcal{R}$ an inner product by the formula

$$\langle r_f, r_h \rangle_{\mathcal{R}} = \langle f, h \rangle_{\oplus_{k \geq 0} \mathcal{E}_k}.$$

$\mathcal{R}$ becomes a Hilbert space and U is a unitary operator from $\oplus_{k \geq 0} \mathcal{E}_k$ onto $\mathcal{R}$. The space $\mathcal{R}$ has the *reproducing property*

$$(4.6) \quad \langle r_f(Z), e \rangle_{\mathcal{E}} = \langle r_f, r_{E(Z)^*e} \rangle_{\mathcal{R}}, \quad f \in \oplus_{k \geq 0} \mathcal{E}_k, \quad e \in \mathcal{E}.$$

We now let $P = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma \in \mathcal{P}_N^0$ take values on $\mathcal{B}_N(\mathcal{E})$ by the the formula:

$$P(Z) = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Z_\sigma, \quad Z \in \mathcal{B}_N(\mathcal{E}).$$

Note that each $f_e = [c_\sigma]_{\sigma \in \mathbb{F}_N^+} \oplus e$, $e \in \mathcal{E}$, belongs to $\oplus_{k \geq 0} \mathcal{E}_k$, so that we deduce from (4.6) that

$$\langle P(Z)e, e' \rangle_{\mathcal{E}} = \langle r_{f_e}, r_{E(Z)^*e'} \rangle_{\mathcal{R}}, \quad e, e' \in \mathcal{E}.$$

All these considerations suggest to introduce the following set as a convenient domain for the evaluation of the orthogonal polynomials on $\mathcal{O}_N^0(\mathcal{A}_2)$. Define

$$\mathcal{G}_N(\mathcal{E}) = \{(W_1 \dots W_N) \in \mathcal{L}(\mathcal{E})^N \mid W_1 W_1^* + \dots + W_{N-1} W_{N-1}^* < \frac{1}{2i}(W_N - W_N^*)\},$$

the *Siegel upper half-space* of $\mathcal{E}$ (see [9], [15]). There is a linear fractional map (*Cayley transform*) connecting $\mathcal{G}_N(\mathcal{E})$ with $\mathcal{B}_N(\mathcal{E})$, defined by the formula:

$$\mathcal{C}(Z) = ((I + Z_N)^{-1} Z_1, \dots, (I + Z_N)^{-1} Z_{N-1}, i(I + Z_N)^{-1}(I - Z_N)),$$

for $Z = (Z_1 \dots Z_N) \in \mathcal{B}_N(\mathcal{E})$. Note that since $(Z_1 \dots Z_N) \in \mathcal{B}_N(\mathcal{E})$, each Z_k , $1 \leq k \leq N$, is a strict contraction ($\|Z_k\| < 1$), hence $\mathcal{C}(Z)$ is well-defined.

Proposition 6. $\mathcal{C}$ is a bijection from $\mathcal{B}_N(\mathcal{E})$ onto $\mathcal{G}_N(\mathcal{E})$.

Proof. For $Z \in \mathcal{B}_N(\mathcal{E})$ we define $W_k = (I + Z_N)^{-1}Z_k$ for $1 \leq k \leq N-1$, and $W_N = i(I + Z_N)^{-1}(I - Z_N)$. Then

$$\begin{aligned} \frac{1}{2i}(W_N - W_N^*) &= (I + Z_N)^{-1}(I - Z_N Z_N^*)(I + Z_N^*)^{-1} \\ &> (I + Z_N)^{-1}(Z_1 Z_1^* + \dots + Z_{N-1} Z_{N-1}^*)(I + Z_N^*)^{-1} \\ &= W_1 W_1^* + \dots + W_{N-1} W_{N-1}^*, \end{aligned}$$

so that $\mathcal{C}(Z) \in \mathcal{G}_N(\mathcal{E})$.

If $W = (W_1 \dots W_N) \in \mathcal{G}_N(\mathcal{E})$, then the imaginary part of W_N is strictly positive, therefore $i + W_N$ is invertible and $\mathcal{C}$ is one-to-one. Also, one easily verifies that for $W \in \mathcal{G}_N(\mathcal{E})$,

$$\mathcal{C}^{-1}(W) = (2i(i + W_N)^{-1}W_1, \dots, 2i(i + W_N)^{-1}W_{N-1}, (i + W_N)^{-1}(i - W_N)).$$

□

We now introduce the Szegő kernel of $\mathcal{G}_N(\mathcal{E})$ by the formula:

$$K_{\mathcal{G}}(W, W') = F(W)F(W')^*, \quad W, W' \in \mathcal{G}_N(\mathcal{E}),$$

where

$$F(W) = 2E(\mathcal{C}^{-1}(W)) \left((i + W_N)^{-1} \right)^{\oplus \infty}.$$

Lemma 7. (a) $K_{\mathcal{G}}$ is a positive definite kernel on $\mathcal{G}_N(\mathcal{E})$.

(b) The set $\{F(W)^* \mathcal{E} \mid W \in \mathcal{G}_N(\mathcal{E})\}$ is total in $\oplus_{k \geq 0} \mathcal{E}_k$.

(c) For any $T \in \mathcal{L}(\mathcal{E})$ and $W, W' \in \mathcal{B}_N(\mathcal{E})$,

$$F(W) \left(\frac{1}{2i}(W_N T - T(W')^*_N) - \sum_{k=1}^{N-1} W_k T(W')^*_k \right)^{\oplus \infty} F(W')^* = T.$$

Proof. (a) and (b) follow directly from Lemma 5. For (c) we notice that if $W = \mathcal{C}(Z)$ and $W' = \mathcal{C}(Z')$ for $Z, Z' \in \mathcal{B}_N(\mathcal{E})$, then

$$T - \sum_{k=1}^N Z_k T(Z')^*_k = 4(i + W_N)^{-1} \left(\frac{1}{2i}(W_N T - T(W')^*_N) - \sum_{k=1}^{N-1} W_k T(W')^*_k \right) (-i + (W')^*_N)^{-1},$$

and together with Lemma 5(c), this concludes the proof. □

We can define

$$\mathcal{S} = \{s_f : \mathcal{G}_N(\mathcal{E}) \rightarrow \mathcal{E} \mid s_f(W) = F(W)f, f \in \oplus_{k \geq 0} \mathcal{E}_k\},$$

and with the inner product

$$\langle s_f, s_h \rangle_{\mathcal{S}} = \langle f, h \rangle_{\oplus_{k \geq 0} \mathcal{E}_k},$$

$\mathcal{S}$ becomes Hilbert space. The connection between the spaces $\mathcal{R}$ and $\mathcal{S}$ is consistent with the connection between the Hardy spaces H^2 and $H^2(\Im z > 0)$ on the unit disk and, respectively, the upper half plane, to which $\mathcal{R}$ and $\mathcal{S}$ reduce for $N = 1$ and $\mathcal{E} = \mathbb{C}$.

We now let $P = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma \in \mathcal{P}_N^0$ take values on $\mathcal{G}_N(\mathcal{E})$ by the formula:

$$P(W) = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma W_\sigma, \quad W \in \mathcal{G}_N(\mathcal{E}).$$

Let ϕ be a strictly positive functional on $\mathcal{O}_N^0$ and let $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ be the associated orthogonal polynomials. Then

$$K_n(W, W') = \sum_{|\sigma| \leq n} \varphi_\sigma(W) \varphi_\sigma(W')^*,$$

is a positive definite kernel on $\mathcal{G}_N(\mathcal{E})$, called the *Christoffel-Darboux kernel*, and the Christoffel-Darboux formula is supposed to provide a connection between K_n and $K_{\mathcal{G}}$.

One of the reasons for the interest in K_n in the classical, one variable case, is that K_n is a reproducing kernel for the set of polynomials of degree at most n with respect to the inner product induced by ϕ . A similar result can be obtained in the non-commutative case. Thus, we have to consider $\mathcal{P}_N^0(\mathcal{E})$, the set of elements of the form $P = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma$, with only finitely many $c_\sigma \in \mathcal{L}(\mathcal{E})$ different from the zero operator on $\mathcal{E}$. On $\mathcal{P}_N^0(\mathcal{E})$ we introduce the product $\sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma \sum_{\tau \in \mathbb{F}_N^+} d_\tau Y_\tau = \sum_{\sigma, \tau \in \mathbb{F}_N^+} c_\sigma d_\tau Y_{\sigma\tau}$ and extend ϕ to $\mathcal{P}_N^0(\mathcal{E})$ by setting $\phi(\sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma) = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma \phi(Y_\sigma)$. Also, extend $\mathcal{I}$ by $\mathcal{I}(\sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma) = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma^* \mathcal{I}(Y_\sigma)$ and define

$$\langle P, Q \rangle_\phi = \phi(\mathcal{I}(Q)P)$$

for $P, Q \in \mathcal{P}_N^0(\mathcal{E})$. Let $P = \sum_{|\sigma| \leq n} c_\sigma Y_\sigma \in \mathcal{P}_N^0$, then $P = \sum_{|\sigma| \leq n} a_\sigma \varphi_\sigma$, where $a_\sigma = \langle P, \varphi_\sigma \rangle_\phi$, so that K_n has the following reproducing property:

$$\begin{aligned} P(W') &= \sum_{|\sigma| \leq n} a_\sigma \varphi_\sigma(W') \\ &= \sum_{|\sigma| \leq n} \langle P, \varphi_\sigma \rangle_\phi \varphi_\sigma(W') \\ &= \langle P, \sum_{|\sigma| \leq n} \varphi_\sigma \varphi_\sigma(W')^* \rangle_\phi \\ &= \langle P, K_{n, W'} \rangle_\phi, \end{aligned}$$

where $K_{n, W'}(W) = K_n(W, W')$ and $W, W' \in \mathcal{G}_N(\mathcal{E})$.

We can obtain now a Cristoffel-Darboux type formula. It is not as simple and relevant as in the commutative case.

Theorem 8. *The Christoffel-Darboux kernel and the Szegő kernel are related by the formula:*

$$\begin{aligned} K_n(W, W') &= F(W) \left(\frac{1}{2i} \left(\Phi_{n+1}(W) B_{n,N} \Phi_n(W')^* - \Phi_n(W) B_{n,N}^* \Phi_{n+1}(W')^* \right) \right) F(W')^* \\ &\quad - \sum_{k=1}^{N-1} F(W) W_k K_n(W, W') (W')_k^* F(W')^*. \end{aligned}$$

Proof. Using the recurrence equations for the orthogonal polynomials we deduce

$$\begin{aligned} W_N K_n(W, W') &- K_n(W, W') (W'_N)^* \\ &= \sum_{k=0}^n \Phi_{k+1}(W) B_{k,N} \Phi_k(W')^* + \sum_{k=0}^n \Phi_{k-1}(W) B_{k-1,N}^* \Phi_k(W')^* \\ &\quad - \sum_{k=0}^n \Phi_k(W) B_{k,N}^* \Phi_{k+1}(W')^* - \sum_{k=0}^n \Phi_k(W) B_{k-1,N} \Phi_{k-1}(W')^* \\ &= \Phi_{n+1}(W) B_{n,N} \Phi_n(W')^* - \Phi_n(W) B_{n,N}^* \Phi_{n+1}(W')^*. \end{aligned}$$

The proof is concluded by an application of Lemma 7 (c). $\square$

5. JACOBI MATRICES

Let ϕ be a strictly positive functional on $\mathcal{O}_N^0(\mathcal{A}_2)$. In addition to the Hilbert space $\mathcal{H}_\phi$, the GNS construction produces a representation of $\mathcal{O}_N^0(\mathcal{A}_2)$ by operators on $\mathcal{H}_\phi$. In this section we analyse this representation in some details by showing the connection with certain matrices of Jacobi type. Let $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ be the family of orthogonal polynomials associated to ϕ . Then we define for $P \in \mathcal{O}_N^0(\mathcal{A}_2) (= \mathcal{P}_N^0)$,

$$(5.1) \quad \Psi_\phi(P) \varphi_\sigma = P \varphi_\sigma.$$

Formula (3.3) shows that each $\Psi_\phi(P)$ is a symmetric operator on $\mathcal{H}_\phi$ with dense domain $\mathcal{D}$, the linear space generated by the polynomials φ_σ , $\sigma \in \mathbb{F}_N^+$. Also, for $P, Q \in \mathcal{P}_N^0$,

$$\Psi_\phi(PQ) = \Psi_\phi(P) \Psi_\phi(Q),$$

and $\Psi_\phi(P) \mathcal{D} \subset \mathcal{D}$ for any $P \in \mathcal{P}_N^0$, hence Ψ_ϕ is an unbounded representation of $\mathcal{O}_N^0(\mathcal{A}_2)$. Also, $\phi(P) = \langle \Psi_\phi(P) 1, 1 \rangle_\phi$ for $P \in \mathcal{P}_N^0$. Of special interest are the operators $\Psi_k = \Psi_\phi(Y_k)$, $k = 1, \dots, N$, since $\Psi_\phi(\sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma) = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma \Psi_{\phi, \sigma}$, where $\Psi_{\phi, \sigma} = \Psi_{i_1} \dots \Psi_{i_k}$ for $\sigma = i_1 \dots i_k$. Since each Ψ_k commutes with the complex conjugation, it follows from von Neumann's theorem (Theorem X.7 in [16]) that each Ψ_k admits self-adjoint extensions.

Let $\{e_1, \dots, e_N\}$ be the standard basis of $\mathbb{C}^N$, then $\{e_{i_1 \dots i_k} = e_{i_1} \otimes \dots \otimes e_{i_k} \mid 1 \leq i_1, \dots, i_k \leq N\}$ is an orthonormal basis of the Fock space $\mathcal{F}(\mathbb{C}^N)$. Let W be the unitary operator from $\mathcal{F}(\mathbb{C}^N)$ onto $\mathcal{H}_\phi$ such that $W(e_\sigma) = \varphi_\sigma$, $\sigma \in \mathbb{F}_N^+$. We see that $W^{-1} \mathcal{D}$ is the linear space $\mathcal{D}_0$ generated by e_σ , $\sigma \in \mathbb{F}_N^+$, so that we can define

$$J_k = W^{-1} \Psi_{\phi, k} W, \quad k = 1, \dots, N,$$

on $\mathcal{D}_0$. Each J_k is a symmetric operator on $\mathcal{D}_0$ and by Theorem 3, the matrix of J_k with respect to the orthonormal basis $\{e_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ is

$$J_k = \begin{bmatrix} A_{0,k} & B_{0,k}^* & 0 & \dots \\ B_{0,k} & A_{1,k} & B_{1,k}^* & \\ 0 & B_{1,k} & A_{2,k} & \ddots \\ \vdots & & \ddots & \ddots \end{bmatrix}.$$

We call $(J_1 \dots J_N)$ a Jacobi N -family on $\mathcal{D}_0$. We can state now the main result of this section about the modeling of Jacobi N -families.

Theorem 9. *Let $(J_1 \dots J_N)$ a Jacobi N -family on $\mathcal{D}_0$ such that $B_n = [B_{n,1} \dots B_{n,N}]$ is an upper triangular invertible matrix. Then there exists a unique strictly positive functional ϕ on $\mathcal{O}_N^0(\mathcal{A}_2)$ such that the map*

$$W(e_\sigma) = \varphi_\sigma, \quad \sigma \in \mathbb{F}_N^+,$$

extends to a unitary operator and

$$J_k = W^{-1} \Psi_{\phi,k} W, \quad k = 1, \dots, N.$$

Proof. Since $B_n, n \geq 0$, are invertible matrices, we can uniquely determine the elements $\varphi_\sigma, \sigma \in \mathbb{F}_N^+$, in $\mathcal{P}_N^0$ such that for $k = 1, \dots, N$,

$$Y_k \varphi_\emptyset = [\varphi_\sigma]_{|\sigma|=1} B_{0,k} + \varphi_\emptyset A_{0,k},$$

and for $k = 1, \dots, N, n \geq 1$,

$$Y_k [\varphi_\sigma]_{|\sigma|=n} = [\varphi_\sigma]_{|\sigma|=n+1}^T B_{n,k} + [\varphi_\sigma]_{|\sigma|=n}^T A_{n,k} + [\varphi_\sigma]_{|\sigma|=n-1}^T B_{n-1,k}^*.$$

Theorem 4 gives a unique strictly positive functional ϕ on $\mathcal{O}_N^0(\mathcal{A}_2)$ such that $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ is the family of orthogonal polynomials associated to ϕ . The GNS construction for this ϕ will produce the required W and $\Psi_{\phi,k}$, as explained above. $\square$

We conclude with an application concerning the moments of positive functionals. Let ϕ be a positive functional on $\mathcal{O}_N^0(\mathcal{A}_2)$. The numbers

$$(5.2) \quad s_\sigma = \phi(Y_\sigma), \quad \sigma \in \mathbb{F}_N^+,$$

are called the *moments* of ϕ . A *Hamburger type problem* would be to determine conditions on the family $\{s_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ so that there exists a positive functional on $\mathcal{O}_N^0(\mathcal{A}_2)$ satisfying (5.2). A solution to this problem can be obtained as follows.

Theorem 10. *The numbers $s_\sigma, \sigma \in \mathbb{F}_N^+$, are the moments of a positive functional on $\mathcal{O}_N^0(\mathcal{A}_2)$ if and only if $K(\sigma, \tau) = s_{I(\sigma)\tau}, \sigma, \tau \in \mathbb{F}_N^+$, is a positive definite kernel on $\mathbb{F}_N^+$.*

Proof. The classical approach extends to this setting. Thus, we notice first that

$$\begin{aligned}
K(\alpha\sigma, \tau) &= s_{I(\alpha\sigma)\tau} \\
(5.3) \qquad &= s_{I(\sigma)I(\alpha)\tau} \\
&= K(\sigma I(\alpha), \tau),
\end{aligned}$$

for $\alpha, \sigma, \tau \in \mathbb{F}_N^+$. Define the sesquilinear form on $\mathcal{P}_N^0$ by the formula

$$\left\langle \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma, \sum_{\tau \in \mathbb{F}_N^+} d_\tau Y_\tau \right\rangle = \sum_{\sigma, \tau \in \mathbb{F}_N^+} s_{I(\sigma)\tau} c_\tau \bar{d}_\sigma.$$

Let $\mathcal{N} = \{P \in \mathcal{P}_N^0 \mid \langle P, P \rangle = 0\}$ and let $\mathcal{H}$ be the Hilbert space obtained by completing $\mathcal{P}_N^0/\mathcal{N}$ in the inner product $\langle \cdot, \cdot \rangle$. Define the maps $\Psi_k : \mathcal{P}_N^0 \rightarrow \mathcal{P}_N^0$, $k = 1, \dots, N$, by

$$\Psi_k \left(\sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma \right) = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_{k\sigma}.$$

From (5.3) we deduce that each Ψ_k , $k = 1, \dots, N$, is a symmetric operator and $\Psi_k \mathcal{N} \subset \mathcal{N}$. Thus, each Ψ_k lifts to a symmetric operator, still denoted Ψ_k , with domain $\mathcal{P}_N^0/\mathcal{N}$ in $\mathcal{H}$. Also $\Psi_k(\mathcal{P}_N^0/\mathcal{N}) \subset \mathcal{P}_N^0/\mathcal{N}$, so that $\Psi_\sigma = \Psi_{i_1} \dots \Psi_{i_k}$ is defined for any $\sigma = i_1 \dots i_k \in \mathbb{F}_N^+$ and we can take

$$\phi \left(\sum_{\sigma \in \mathbb{F}_N^+} c_\sigma Y_\sigma \right) = \left\langle \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma \Psi_\sigma \hat{1}, \hat{1} \right\rangle,$$

where $\hat{1}$ denotes the class of $1 \in \mathcal{P}_N^0$ in $\mathcal{P}_N^0/\mathcal{N}$. One can easily check that ϕ is a solution of the Hamburger problem with data $\{s_\sigma\}_{\sigma \in \mathbb{F}_N^+}$. $\square$

A more specialized version of this result was proved in [10].

REFERENCES

- [1] W. Arveson, Subalgebras of C^* -algebras III: Multivariable operator theory, *Acta Math.*, **181**(1998), 476–514.
- [2] C. Berg, J. P. R. Christensen, and P. Ressel, *Harmonic Analysis on Semigroups*, Springer, Berlin, 1984.
- [3] K. C. Chou, A. S. Wilsky, and A. Benveniste, Multiscale recursive estimation, data fusion and regularization, *IEEE Trans. Automatic Control*, **39**(1994), 464–478.
- [4] T. Constantinescu, Schur analysis of positive block matrices, in *Schur Methods in Operator Theory and Signal Processing*, (I. Gohberg, Ed.), pp. 191–206, Birkhäuser, Basel, 1986.
- [5] T. Constantinescu, *Schur Parameters, Factorization and Dilation Problems*, Birkhäuser, Basel, 1996.
- [6] T. Constantinescu and J. L. Johnson, A note on noncommutative interpolation, *Canad. Math. Bull.*, to appear.
- [7] T. Constantinescu and J. L. Johnson, Tensor algebras and displacement structure. II. Noncommutative Szegő theory, preprint 2002.
- [8] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher transcendental functions*, McGraw-Hill, New York, 1953.

- [9] G. B. Folland, *Harmonic Analysis in Phase Space*, Annals of Math. Studies, Princeton Univ. Press, 1989.
- [10] D. Hadwin, A noncommutative moment problem, *Proc. Amer. Math. Soc.*, **129**(2001), 1785–1791.
- [11] T. H. Koornwinder, Special functions and q -commuting variables, in *Special Functions, q -Series and Related Topics*, (M. E. H. Ismail, D. R. Masson, M. Rahman, Eds.), pp. 131–166, Fields Institute Communications, Amer. Math. Soc., Providence, Rhode Island, 1997.
- [12] M. A. Kowalski, The recursion formulas for orthogonal polynomials in n variables, *SIAM J. Math. Anal.*, **13**(1982), 309–323.
- [13] E. C. Lance, *Hilbert C^* -modules. A toolkit for operator algebraists*, Cambridge Univ. Press, Cambridge, 1995.
- [14] K. R. Parthasarathy, *An Introduction to Quantum Stochastic Calculus*, Birkhäuser, Basel 1992.
- [15] R. M. Range, *Holomorphic Functions and Integral Representations in Several Complex Variables*, Springer, Berlin, 1986.
- [16] M. Reed and B. Simon, *Methods of Modern Mathematical Physics*, II., Academic Press, New York, 1975.
- [17] G. Szegő, *Orthogonal polynomials*, Colloquium Publications, **23**, Amer. Math. Soc., Providence, Rhode Island, 1939.
- [18] Y. Xu, On orthogonal polynomials in several variables, in *Special Functions, q -Series and Related Topics*, (M. E. H. Ismail, D. R. Masson, M. Rahman, Eds.), pp. 247–270, Fields Institute Communications, Amer. Math. Soc., Providence, Rhode Island, 1997.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT DALLAS, BOX 830688, RICHARDSON, TX 75083-0688, U. S. A.

E-mail address: `tiberiu@utdallas.edu`